

1115 a. *Eucereon relegata*. (Plate XIX. fig. 15.)

*Eucereon relegatum*, Schaus, A. M. N. H. (8) vii. p. 182 (1911).

♂. Head and thorax grey-brown suffused and obscurely streaked with black, the back of head with two slight yellow spots; palpi with the first joint ochreous in front; coxæ, streaks on femora, and rings at end of tibiæ and base of tarsi and spurs ochreous; abdomen orange-yellow with black-brown triangular patch on basal half of dorsum followed by a black fascia, the extremity and lateral fascia black, the ventral surface ochreous white except terminal segments. Fore wing ochreous mostly suffused with fuscous brown, the veins with dark streaks; subbasal black spots defined by ochreous below costa and cell; an antemedial series of four elongate spots followed by four other spots, the spot in cell large and the three below the cell obliquely placed; a wedge-shaped spot in upper angle of cell and elongate spots in and below lower angle; four spots beyond the cell; an obliquely curved postmedial series of nine elongate spots; a subapical spot and six terminal spots. Hind wing semi-hyaline white, the veins and margins fuscous, the fuscous on termen rather diffused and angled inwards below vein 2.

*Hab.* COSTA RICA, Juan Vinas (*Schaus*), 1 ♂. *Exp.* 42 millim.