

1112 *b.* **Eucereon consorta.** (Plate XIX. fig. 12.)

Eucereon consorta, Schaus, A. M. N. H. (8) vi. p. 192 (1910).

Head and thorax white; palpi black, the 2nd joint white in front towards tips, the 3rd joint white at tips; frons with the lower part black at sides; antennæ black; tegulæ edged with black; basal half of patagia with two black streaks; metathorax with black patch; pectus and legs black, the coxæ white, the femora and mid and hind tibiæ with white spots; abdomen yellow with large dorsal black-brown patch extending to 6th segment, the extremity black, lateral series of black patches, the ventral surface white, the terminal segments black. Fore wing ochreous white; an irregular black patch on costa near base, produced in cell and connected with a black fascia on basal part of inner margin expanding into two projections above vein 1 before middle; a small antemedial black spot below the cell and two confluent spots on vein 2 near its base; a round spot in middle of cell connected with a patch on costa; a large irregular patch from costa to lower angle of cell; a patch formed of three conjoined marks, the upper one minute, from below vein 2 to inner margin and an oblique triangular mark with minute spot above its outer angle from below end of cell to termen just above tornus; a patch from costa towards apex to discal fold, its outer edge angled outwards below a spot on costa just before apex; a spot on termen below vein 7, point at discal fold, and spots below veins 5 and 4. Hind wing fuscous, the lower part of cell and areas just below and beyond it semihyaline, especially in male.

Hab. COSTA RICA, Tuis (*Schaus*), 1 ♂, 2 ♀, Juan Vinas (*Schaus*), 3 ♀, Sitio (*Schaus*), 1 ♀. *Exp.* 46 millim.