

b. Antennæ of male serrate and fasciculate.

a¹. Fore wing of male with small glandular swelling and tuft of hair at base of costa on underside; hind wing with the costa fringed with long hair.

1249. *Automolis herois*.

Idalus herois, Schaus, Ent. Am. v. p. 190 (1889); Druce, Biol. Centr.-Am., Het. ii. p. 366, pl. 73. f. 20; Kirby, Cat. Het. p. 198.

♂. Head and thorax white; antennæ brownish; palpi and frons fuscous; vertex of head with orange spot; tegulæ and patagia with orange spots, the tegulæ tipped with orange, the patagia edged with crimson; legs with fuscous points; abdomen crimson above, white below. Fore wing pale yellow, with the inner area and the medial part of costa white; the basal area with blue-black




Fig. 34.—*Automolis herois*, ♂. $\frac{1}{2}$.

streaks, not reaching the base on inner area; a medial blue-black band, produced outwards on costa and inner margin and with the veins on it whitish. Hind wing with the inner half tinged with crimson.

Hab. MEXICO, Coatepec (*Brooks*), 2 ♂, Godman-Salvin Coll.; VENEZUELA, Merida; BRAZIL, Sta. Catherina. *Exp.* 38 millim. Type in Coll. Schaus.