

*1124 *a.* **Eucereon pseudocasca.** (Plate XIX. fig. 29.)

Eucereon pseudocasca, Roths. Nov. Zool. xix. p. 174 (1912).

♂. Head black with two yellow spots behind; palpi with some yellow at base; thorax blackish and grey, the patagia edged with yellow, the metathorax with white patch; pectus and legs black, the femora and tibiae streaked with whitish; abdomen black, the anal tuft at base and sides yellow, the basal half of ventral surface white. Fore wing whitish, the costa and veins beyond the cell streaked with yellow; the subbasal area black-brown from costa to above inner margin, interrupted by pale streaks on the veins and in submedian fold; an antemedial series of slight black-brown marks from costa to median nervure; the medial area with black-brown streaks below costa, above and below basal half of vein 2, and from submedian fold to inner margin; a large lunulate black-brown discoidal patch extending from costa to below end of cell, broken up by the yellow streaks on the veins; two postmedial spots above inner margin; the apical area black-brown to below vein 3, broken up by the yellow streaks on the veins and traversed by the whitish subterminal line which is angled outwards at vein 6 and bent outwards to termen at submedian fold; two black-brown subterminal spots above tornus. Hind wing semihyaline whitish, the inner area tinged with brown, the terminal area fuscous brown, broad at costa, narrowing to tornus, its inner edge irregular; a dark mark on upper discocellular; the underside with the costal area suffused with fuscous brown.

♀. Abdomen with the extremity of penultimate and base of anal segment yellow.

Hab. PERU, Carabaya, S. Domingo (*Ockenden*), type † ♂, ♀ in Coll. Rothschild. *Exp.* 36 millim.