

1094. **Eucereon æolum**, n. sp. (Plate XVI. fig. 16.)

Eucereon rhodophila, Druce, Biol. Centr.-Am., Het. i. p. 86 (nec Wlk.).

Pale brown; head, tegulæ, and thorax spotted with black; back of head and metathorax with orange spots; patagia streaked with black; legs streaked with yellowish grey; abdomen crimson, with dorsal black-brown patches decreasing to 3rd segment and followed by a series of points, the terminal segment black, lateral black fasciæ, the ventral surface greyish yellow. Fore wing pale brown; basal and subbasal series of short streaks, two of the latter series meeting on vein 1; an antemedial series of black spots followed by two spots in cell; two spots in end of cell, and a series just beyond the cell bent inwards below the cell to near the antemedial series; a postmedial series of spots oblique below vein 4, and followed by a series of dentate marks from costa to vein 4; a subterminal series of spots and a terminal series of points on the veins. Hind wing pale semihyaline, the veins, inner and terminal areas suffused with fuscous; veins 3, 4, 5 from angle.

Hab. MEXICO, Cordova (*Rimeli*), type, Coatepec (*Brooks*), Tabasco (*H. H. Smith*), Jalapa (*Trujillo*); GUATEMALA (*Champion*): Godman-Salvin Coll. *Exp.* 34 millim.