

1121 *e.* **Eucereon zephyrus.** (Plate XIX. fig. 25.)

Eucereon zephyrum, Schaus, A. M. N. H. (8) vi. p. 194 (1910).

♂. Head white, the frons with slight fuscous spots at middle and above, the back of head with pair of yellowish white patches; palpi black except at base; antennæ black; thorax white mixed with greyish fuscous; legs blackish streaked with white; abdomen pale yellow, greyish at base, then with fuscous dorsal streak, the anal tuft grey and fuscous, the ventral surface white and blackish. Fore wing white, the veins, discal and submedian folds streaked with fuscous, the basal half of costal edge blackish; a fuscous spot above vein 1 before the antemedial line, which is fuscous, strongly excurved in cell and slightly below submedian fold; a slight fuscous discoidal lunule; postmedial line fuscous, slightly angled outwards below costa, excurved at middle, then strongly bent inwards and angled outwards at vein 1; subterminal line represented by a diffused fuscous band from costa to vein 5 and a band from vein 3 to inner margin bent inwards at vein 2; a fuscous line just

Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: 347

before termen from costa to below vein 5, then a blackish bar on termen to vein 3. Hind wing semihyaline white, the apical area suffused with fuscous, the termen fuscous.

Hab. COSTA RICA, Juan Vinas (*Schaus*), 1 ♂. *Exp.*
34 millim.

E. zephyrum *Schaus.* Anterior body grey-white, hindhead and neck like the abdomen above yellow, *zephyrum*. the latter above with subdorsal grey stripes, the two last segments all grey, beneath grey. The white forewings have grey veins and stripes on the cellular and submedian folds; anterior transverse line dentate; the posterior one, encircling the large discal spot, runs from the upper median vein to the middle of the proximal margin; behind it a large, white area with likewise white veins, in the apex and at the anal angle a greyish-spotted obscuration. The hindwings semi-transparent, iridescent, with a narrow, dark border. From Costa Rica. — **songoense** *subsp. nov.* (25 h) has above a quite unicolorous grey abdomen, being all white beneath; *songoense*. the cellular and submedian folds are not striped grey, but the veins are grey-brown also in the marginal area, at any rate all the markings brownish. Bolivia (Rio Songo), from BANG-HAAS. Draudt, 1915; Gross-Schmett. Erde 6: 183