

1131. Eucereon discolor.

Eucerea discolor, Wlk. vii. 1640 (1856); Kirby, Cat. Het. p. 201.

Eucereon rububa, Druce, A. M. N. H. (6) xv. p. 46 (1895); id. Biol. Centr.-Am., Het. ii. p. 364, pl. 73. f. 17.

♂. Head black-brown; frons with metallic blue patch; thorax and base of abdomen greyish brown and black, the rest of abdomen black, with pale ventral patches on first two segments. Fore wing blackish brown, the inner area pale, conjoined to diffused pale anter- and postmedial bands; a pale spot in end of cell; obliquely curved obscure dark antemedial, postmedial, and subterminal lines; three diffused pale spots on termen. Hind wing pale, the apical area and termen suffused with fuscous.

Fore wing with vein 6 from below angle of cell; hind wing with vein 3 separate from 4, 5.

Hab. COSTA RICA (*Underwood*), 1 ♂, type *rububa*, Godman-Salvin Coll.; PANAMA; BRAZIL, São Paulo (*Jones*), 2 ♀, Rio Janeiro. Exp. 30–46 millim. Type † in Mus. Oxon.

Subsp. 1. Geyser; hind wing entirely white except apical area.
Hab. BOLIVIA, R. Tanampaya (*Garlepp*), 1 ♀. Exp. 46 millim.