*1102 d. Eucereon griseata. (Plate XIX. fig. 5.)

Eucereon griseatum, Roths. Nov. Zool. xix. p. 174 (1912).

Q. Head black with two orange spots behind; palpi with white spots at extremity of 2nd joint in front; frons with white bar above; antennæ with the shaft ringed with white towards base; thorax black, the tegulæ with white spots at sides, the shoulders with orange spots, the patagia white edged with black and with streak at middle, the dorsum of thorax with pair of white streaks; pectus white at sides, the femora and tibiæ streaked with white; abdomen black-brown with quadrate orange patch on two terminal segments and sublateral series of whitish patches except towards extremity. Fore wing creamy white suffused with olive-yellow to the postmedial line, the veins, discal and submedian folds, and inner margin streaked with black; antemedial line black, forming a spot at costa, bent outwards in upper part of cell, incurved at median nervure, and angled outwards in submedian fold; a spot on costa above end of cell with an incurved line from it to the

Hampson, 1914, Cat. Lepid. Phalaenae Br. Mus. (Suppl.) 1: 335

336

AMATIDE.

discocellulars at discal fold; postmedial line black, arising from the spot above end of cell, oblique and sinuous to vein 3, then bent inwards to vein 2 near its origin and incurved to inner margin; subterminal line black, rather diffused and waved, excurved and forming a patch below vein 7, angled inwards at vein 5, excurved and connected with the termen at middle, then incurved, forming a patch below vein 3 connected with a spot on termen in submedian fold; a black spot just before apex and terminal line from below apex to vein 4. Hind wing semihyaline brownish white, the terminal area suffused with fuscous narrowing to tornus. Underside of fore wing suffused with fuscous leaving a white patch in cell and oblique maculate band from below costa beyond middle to vein 3 near termen; hind wing with the costal area suffused with fuscous.

Ab. 1. Hind wing wholly suffused with fuscous brown.

Hab. Peru, Carabaya, Tinguri (Ockenden), type † ♀ in Coll. Rothschild, S. Domingo, R. Suambari, La Oroya, Chanchamayo. Exp. 46 mlllim.