

Genus **HYPOCLADIA**, nov.

Type, *H. militaris*.

Proboscis well developed; palpi upturned, the 2nd joint reaching vertex of head, the 3rd moderate; antennæ bipectinate, the branches dilated and ending in a bristle; tibiæ with the spurs moderate. Fore wing rather narrow; vein 3 from near angle of cell; 4, 5 from angle; 6 from upper angle; 7, 8, 9, 10 stalked; 11 from cell. Hind wing with the cell long; vein 2 from near angle; 3, 4 from angle; 5 from above angle; 6, 7 from upper angle.

800. **Hypocladia militaris**.

Dycladia militaris, Butl. Ill. Het. i. p. 33, pl. 16. f. 1 (1877); Kirby, Cat. Het. p. 145.

♂. Head, thorax, and abdomen brown; two spots on vertex of head, on tegulæ and metathorax crimson; coxæ mostly whitish; abdomen with lateral crimson fasciæ decreasing and becoming maculate distally; ventral white bands to near extremity. Fore wing brown, with whitish streaks in and below cell, and a series of short streaks between veins 2 and 7, becoming shorter and maculate towards vein 7. Hind




Fig. 173.

Hypocladia militaris, ♂. $\frac{1}{1}$.

wing black-brown, with slight hyaline streaks in, below, and beyond cell.

Hab. AMAZONS, 1 ♂ type. *Exp.* 28 millim.