

## Genus CLOËSIA, nov.

Type, *C. parthia*.

Proboscis fully developed; palpi upturned, not reaching vertex of head, the 2nd joint fringed with hair in front; antennæ of male with bristles and cilia; tibiæ with the spurs short. Fore wing with vein 2 from towards angle of cell; 3 from before angle; 4, 5 from angle; 6 from below upper angle; 7, 8, 9 stalked; 10 from cell; 11 anastomosing with 12. Hind wing with vein 2 from beyond middle of cell; 3 from close to angle; 5 from well above angle; 6, 7 stalked; 8 from middle of cell and strongly curved.

### 727. *Cloësia parthia*.

*Ichoria parthia*, Druce, A. M. N. H. (6) iv. p. 83 (1889); id. Biol. Centr. Am., Het. ii. p. 336, pl. 70. f. 23; Kirby, Cat. Het. p. 128.




Fig. 252.—*Cloësia parthia*, ♂. 1.

Head and thorax deep metallic blue-green; abdomen deep purplish blue. Fore wing metallic golden green. Hind wing blue-green.

*Hab.* NICARAGUA, Chontales (*Belt*), 1 ♂; PANAMA, Bugaba (*Champion*), 1 ♂, type, Godman-Salvin Coll.; BRAZIL, São Paulo. *Exp.* 34 millim.