

83. *Automolis pseudameoides* spec. nov.

♂. This species resembles *Prumala ameoides*. Palpi and legs brownish black, streaked longitudinally with yellowish white; head and collar white; tegulae golden yellow, broadly edged on inner side with black; patagia golden yellow; thorax golden yellow, with central black line; abdomen velvety black, yellow on sides of first two segments, and having white streaks on last four segments.—Forewing: costal edge and vein 11 buff, rest of wing black; from base to angle of inner margin placed obliquely is a large diamond-shaped golden yellow patch occupying one-third of the wing area; in the apical half of the wing is another large golden yellow irregularly triangular patch, and between the two from the outer margin to edge of cell runs a narrow buff line.—Hindwing yellowish buff, with a broad band of sooty brownish reaching from anal angle to middle of outer margin; costal third of the wing occupied by an enormous scent gland and patch of androconia.

♀. Similar, but sooty brown; abdomen above and on sides entirely deep brown glossed with blue, and merely a few whitish scales near tip; the two golden yellow patches on forewing smaller; veins whitish.—Hindwing deep sooty brown; basal third yellow.

Length of forewing: ♂ 19 mm.; ♀ 21 mm.

*Hab.* La Union, R. Huacamayo, Carabaya, Peru, November 1904 (G. Ockenden).

7 ♂♂, 2 ♀♀.