

1249 c. *Automolis ochreata*.

Automolis ochreata, Schaus, Pr. U.S. Nat. Mus.*xxix. p. 214 (1905);
Roths. Nov. Zool. xvi. pl. 5. f. 30.

Fore wing of male without swelling and tuft of hair at base of costa; large elliptical patches of dark brown scales below middle of cell of fore wing on underside and below costa of hind wing on




Fig. 58.—*Automolis ochreata*, ♂. $\frac{1}{2}$.

upper side; patches of white scales on fore wing above middle of vein 1 on underside and on hind wing in and beyond end of cell on upper side; hind wing with veins 2, 3 from a point, 5 from above middle of discocellulars.

♂. Head yellow, 1st joint of palpi and frons white, 2nd and 3rd joints of palpi brown; tegulae and shoulders white with blackish spots at sides; thorax yellow, the metathorax white with pair of black points; pectus and legs yellow and white; abdomen yellow, dorsally tinged with crimson on basal half. Fore wing orange-yellow; an oblique white antemedial band with series of black spots on it, rather constricted at middle and conjoined at inner margin to an erect postmedial white band with series of elongate black marks on it in the interspaces, those at end of cell bisected by white on the discocellulars. Hind wing yellow, the inner half tinged with crimson and with deeper crimson streak below the cell.

Hab. FR. GUIANA, Maroni R. (*Schaus*), 1 ♂. *Exp.* 32 millim.