

N. clara *Cr.* (149 k) is easily recognisable, very different from the other species, the brown forewing *elara*. with black longitudinal stripes and a large triangular silvery white spot in the disc. Guiana, Colombia.

N. marchiana *Schs.* is extremely similar to *clara*, discernible by the antennae being tuftedly ciliated *marchiana*. in the basal halves, not shortly pectinated. The broad black subterminal line of the forewing forms internerval lunules, only one between 6 and 4, behind it follows a fine red-brown lunular line, interrupted on the veins. Hindwing whitish, brownish sand-coloured at the margins, narrowly so at the distal margin. Expanse of wings: 40 mm. Colombia (Eastern Andes); Bolivia (Rio Songo). Genitally very different from *clara*.

Draudt in Seitz

N. eugenia *Schs.* Basal part of forewing, from the first quarter of the costal margin to the centre of the *eugenia*. inner margin, dark brown with velvety darker stripes, reddish-yellow behind it, lilac at the costal margin, strewn with brown below it, cell-end whitish, with an oval brown line; the posterior transverse line is fine, dark brown, shaded with brown and grey behind it, with velvety darker internerval streaks; undulate line whitish, widened between the veins 4 and 6. Hindwing dark grey. Expanse of wings: 27 mm. French Guiana.